ΑD			

Award Number:

W81XWH-07-1-0374

TITLE:

Mesenchymal Stem Cell as Targeted-Delivery Vehicle in Breast Cancer

PRINCIPAL INVESTIGATOR:

Zhen Cheng

CONTRACTING ORGANIZATION:

Stanford University School of Medicine

Stanford, CA 94305-5484

REPORT DATE:

June 2010

TYPE OF REPORT:

Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT:

 $\Box x$ Approved for public release; distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE					Form Approved OMB No. 0704-0188		
data needed, and completing this burden to Department of I 4302. Respondents should be	and reviewing this collection of in Defense, Washington Headquar The aware that notwithstanding an	nformation. Send comments regiters Services, Directorate for Info	arding this burden estimate or an rmation Operations and Reports n shall be subject to any penalty	y other aspect of this co (0704-0188), 1215 Jeffe	thing existing data sources, gathering and maintaining the ollection of information, including suggestions for reducing erson Davis Highway, Suite 1204, Arlington, VA 22202- on a collection of information if it does not display a currently		
1. REPORT DATE (DE 01-06-2010	D-MM-YYYY)	2. REPORT TYPE Annual	NE33.		MAY 2009 - 14 MAY 2010		
4. TITLE AND SUBTIT	rle '			5a.	CONTRACT NUMBER		
Mesenchymal S	tem Cell as Targ	geted-Delivery Ve	hicle in Breast Ca	5b.	GRANT NUMBER 1XWH-07-1-0374		
				5c.	PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)				5d.	PROJECT NUMBER		
Zhen Cheng				5e.	TASK NUMBER		
				5f. \	WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Stanford University School of Medicine, Stanford, CA				8. PERFORMING ORGANIZATION REPORT NUMBER			
9. SPONSORING / MO	ONITORING AGENCY	NAME(S) AND ADDRES	S(ES)	10.	SPONSOR/MONITOR'S ACRONYM(S)		
•	al Research and Ma	ateriel Command					
Fort Detrick, MD	21702-5012				SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION / A	AVAILABILITY STATE	MENT					
Approved for publ	lic release; distribut	ion unlimited					
13. SUPPLEMENTAR	Y NOTES						
agents in tumors. In and dual reporter ge imaging techniques vitro; 3) We will de Major findings from premature and well- potential at lung tun	this proposal we will one mrfp-ttk. Specific (2) We will demonst termine the effect of a year 1 studies: 1) M established breast ca	I stably transfect mese Aims: 1) We will morate the ability of mese mesenchymal stem ce (SCs home to both sub ncer lung metastasis; as tumor niches; 5) lung	enchymal stem cells we nitor breast cancer troe enchymal stem cells to a ll to target delivery of cutaneous breast cance. MSCs proliferate at	ith a lentiviral v pism of mesence target delivery gene therapeuti er and its lung r tumor site; 4) N	and local production of biological rector containing a therapeutic gene hymal stem cells by multimodality of gene therapeutics to breast cancer in cs to breast cancer lung metastasis. metastasis; 2) MSCs home to both MSCs show dissimilar differentiation is BMP-2 and Noggin transcription,		
15. SUBJECT TERMS	1						
oobobo i Ekimo	•						
16. SECURITY CLASSIFICATION OF:		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC			
a. REPORT	b. ABSTRACT	c. THIS PAGE	UU	7	19b. TELEPHONE NUMBER (include area code)		
11	1 11	1 11	1	1			

Table of Contents

<u>Page</u>

Introduction	4
Body	5
Key Research Accomplishments	7
Reportable Outcomes	7
Conclusion	7
References	7
Appendices	7

INTRODUCTION


In our previous annual report, we found that mesenchymal stem cells (MSCs) homing to breast cancer lung metastasis differentiate into osteoblasts, whereas MSCs homing to subcutaneous breast cancer differentiate into adipocytes. The data provide evidence that MSCs home and engraft to tumor site, and that different tissue tumor niche have distinct effect on MSCs differentiation. These finding may be clinically relevant because the beneficial effects of MSCs are being tested clinically in attempts to improve hematopoietic engraftment [1], to treat osteogenesis imperfect [2], graft-versus-host disease [3], and autoimmune diseases [4, 5], and to deliver therapy for malignancies [6, 7].

For the current funding period, we have focused on our specific aim II presented in the research proposal: We will demonstrate the ability of mesenchymal stem cells to target delivery of RGD4CrmhTNF fusion protein to breast cancer cells and endothelial cells in vitro.

BODY

To achieve our goal, we obtained the plasmid RGD-rmhTNF (recombinant mutated human TNF) from our collaborator, Dr. Yinqi Zhang, MD, PhD, Professor and Director of Center of Biotechnology at The Fourth Military Medical University in Xian, China. Then we finished the design of vector and our cloning strategy is listed below. We are actively working on the cloning the vector and expect to obtain the vector very soon.


The lentivirus vector, pLVX-IRES-ZsGreen1 from Clontech, will be used to stably express the fusion protein of rmhTNF-mrfp-ttk stem cells. First, the restriction site of AgeI will be inserted upstream of ZsGreen1 sequence by using GeneTailor Site-Directed Mutagenesis system from Invitrogen. Also, a PacI site will be added at the 3' end of ZsGreen1 sequence by using the same strategy. The fragment of RGDrmhTNF (recombinant mutated human TNF) was amplified from PBV220-RGD-rmhTNF using the sense



primer of 5'-GCGAATTC ATGCGTCGTCGTCGTTCTTCTTATTCTTGTTATGTGC-3' and the antisense primer of 5'- GGAGCGGCCGCTCAGAATTCAATGA-3'. The restriction site of EcoRI and NotI underlined in the primer sequences were added respectively to the 5' and 3' end by PCR. The italicized sequence, ATGCGTCGTCGTCTCTTCTTATTCTTGTT, encodes the signal peptide of MRRRSLLILV. Both PCR product and the modified vector of pLVX-IRES-ZsGreen1 (pLVX-IRES-A-ZsGreen1-P) will be digested with EcoRI and NotI and ligated by T4 ligase. The fragment of mrfp-ttk were amplified from the vector of pcDNA3-hrl-mrfp-ttk (a kind gift from Dr. Gambhir at Stanford University) by using the primer set of 5'-GCACCGGTATGGCCTCCTCCGAGGAC-3' and 5'-ACGCTTAATTAA TCAGTTAGCCTCCCC-3' with AgeI added to the 5' end and PacI inserted to the 3' end, both sites were underlined in the primer sequences. Both PCR product and pLVX-IRES-A-ZsGreen1-P will be digested with AgeI and PacI. The fragment of mrfp-ttk will then be inserted downstream of IRES and replace ZsGreen1 sequence.

In the meanwhile, to choose a TNF sensitive breast cancer cell line, we did 24h MTT assay using several breast cancer cell lines and found MDA-MB-231 is the most sensitive breast cancer cell line among others such as: MDA-MB-435, MDA-MB-468 and MCF-7. Figure 2 is the result of MTT assay.

Based on the MTT assay, we will select MDA-MB-231 cells for in vitro study and for preparation of lung metastasis tumor model. To monitor metastasis tumor growth, we need to transfect reporter gene of firefly luciferase into cells in order to apply bioluminescence imaging. We already did the transfection and cells are currently in the process of antibiotic screening for obtaining stable cell lines.



KEY RESEARCH ACCOMPLISHMENTS

There is no key research accomplishment in this reporting period.

REPORTABLE OUTCOMES

Because of limited progress of the project, there have been no manuscripts, abstracts, or products, patents, and career developments that resulted from this award during the reporting year.

CONCLUSIONS

In conclusion, because of funding transferring process, we lost quite significant amount of research time. The pLVX-IRES-R4CT-mrfp-ttk vector has been under cloning.

REFERENCES

- 1. Lazarus HM, Koc ON, Devine SM, Curtin P, Maziarz RT, Holland HK, et al. Cotransplantation of HLA-identical sibling culture-expanded mesenchymal stem cells and hematopoietic stem cells in hematologic malignancy patients. Biol Blood Marrow Transplant. 2005;11:389-98.
- 2. Horwitz EM, Prockop DJ, Fitzpatrick LA, Koo WW, Gordon PL, Neel M, et al. Transplantability and therapeutic effects of bone marrow-derived mesenchymal cells in children with osteogenesis imperfecta. Nat Med. 1999;5:309-13.
- 3. Le Blanc K, Rasmusson I, Sundberg B, Gotherstrom C, Hassan M, Uzunel M, et al. Treatment of severe acute graft-versus-host disease with third party haploidentical mesenchymal stem cells. Lancet. 2004;363:1439-41.
- 4. Jones OY, Good RA, Cahill RA. Nonmyeloablative allogeneic bone marrow transplantation for treatment of childhood overlap syndrome and small vessel vasculitis. Bone Marrow Transplant. 2004;33:1061-3.
- 5. El-Badri NS, Maheshwari A, Sanberg PR. Mesenchymal stem cells in autoimmune disease. Stem Cells Dev. 2004;13:463-72.
- 6. Studeny M, Marini FC, Champlin RE, Zompetta C, Fidler IJ, Andreeff M. Bone marrow-derived mesenchymal stem cells as vehicles for interferon-beta delivery into tumors. Cancer Res. 2002;62:3603-8.
- 7. Studeny M, Marini FC, Dembinski JL, Zompetta C, Cabreira-Hansen M, Bekele BN, et al. Mesenchymal stem cells: potential precursors for tumor stroma and targeted delivery vehicles for anticancer agents. J Natl Cancer Inst. 2004;96:1593-603.

APPENDICES

None.